arXiv:1104.5420v1 [math.NT] 28 Apr 2011

A note on the generalized q-Bernoulli measures with weight α

D. V. Dolgy, T. Kim, S. H. Lee, C. S. Ryoo

Institute of Mathematics and Computer Science,
Far Eastern National University, Vladivostok, 690060, Russia

Division of General Education,
Kwangwoon University, Seoul 139-701, Korea

Department of Mathematics,
Hannam University, Daejeon 306-791, Korea

Abstract In this paper we discuss new concept of the q-extension of Bernoulli measure. From those measures, we derive some interesting properties on the generalized q-Bernoulli numbers with weight α attached to χ.

2000 Mathematics Subject Classification - 11B68, 11S40, 11S80

Key words- Bernoulli numbers and polynomials, q- Bernoulli numbers and polynomials, q-Bernoulli numbers and polynomials

1. Introduction

Let p be a fixed prime number. Throughout this paper $\mathbb{Z}_p$, $\mathbb{Q}_p$, and $\mathbb{C}_p$ will, respectively, denote the ring of p-adic rational integers, the field of p-adic rational numbers, and the completion of algebraic closure of $\mathbb{Q}_p$. Let $\mathbb{N}$

be the set of natural numbers and $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. Let ν_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-\nu_p(p)} = \frac{1}{p}$ (see [1-14]).

When we talk of q-extension, q is variously considered as an indeterminate, a complex number $q \in \mathbb{C}$, or a p-adic number $q \in \mathbb{C}_p$. Throughout this paper we assume that $q \in \mathbb{C}_p$ with $|1-q|_p < 1$ and we use the notation of q-number as

$$[x]_q = \frac{1-q^x}{1-q}, \text{ (see [1-14]).}$$

Thus, we note that $\lim_{q \to 1}[x]_q = x$.

In [2], Carlitz defined a set of numbers $\xi_k = \xi_k(q)$ inductively by

$$\xi_0 = 1, \quad (q\xi+1)^k - \xi_k = \begin{cases} 1, & \text{if } k=1, \\ 0, & \text{if } k>1, \end{cases} \tag{1}$$

with the usual convention of replacing ξ^k by ξ_k.

These numbers are q-extension of ordinary Bernoulli numbers B_k. But they do not remain finite when $q=1$. So he modified (1) as follows:

$$\beta_{0,q} = 1, \quad q(q\beta+1)^k - \beta_{k,q} = \begin{cases} 1, & \text{if } k=1, \\ 0, & \text{if } k>1, \end{cases} \tag{2}$$

with the usual convention of replacing β^k by $\beta_{k,q}$.

The numbers $\beta_{k,q}$ are called the k-th Carlitz q-Bernoulli numbers.

In [1], Carlitz also considered the extended Carlitz's q-Bernoulli numbers as follows:

$$\beta^h_{0,q} = \frac{h}{[h]_q}, \quad q^h(q\beta^h+1)^k - \beta^h_{k,q} = \begin{cases} 1, & \text{if } k=1, \\ 0, & \text{if } k>1, \end{cases}$$

with the usual convention of replacing $(\beta^h)^k$ by $\beta^h_{k,q}$.

Recently, Kim considered q-Bernoulli numbers, which are different extended Carlitz's q-Bernoulli numbers, as follows: for $\alpha \in \mathbb{N}$ and $n \in \mathbb{Z}_+$,

$$\widetilde{\beta}^{(\alpha)}_{0,q} = 1, \quad q(q^\alpha \widetilde{\beta}^{(\alpha)}+1)^n - \widetilde{\beta}^{(\alpha)}_{n,q} = \begin{cases} \frac{\alpha}{[\alpha]_q}, & \text{if } n=1, \\ \\ 0, & \text{if } n>1, \end{cases} \tag{3}$$

with the usual convention of replacing $(\widetilde{\beta}^{(\alpha)})^k$ by $\widetilde{\beta}^{(\alpha)}_{k,q}$ (see [3]).

The numbers $\widetilde{\beta}_{k,q}^{(\alpha)}$ are called the k-th q-Bernoulli numbers with weight α.

For fixed $d \in \mathbb{Z}_+$ with $(p,d)=1$, we set

$$\begin{aligned} X &= X_d = \varprojlim_N (\mathbb{Z}/dp^N\mathbb{Z}), \quad X_1 = \mathbb{Z}_p, \\ X^* &= \bigcup_{\substack{0<a<dp \\ (a,p)=1}} (a + dp\mathbb{Z}_p), \\ a + dp^N\mathbb{Z}_p &= \{x \in X \mid x \equiv a \pmod{dp^N}\}, \end{aligned}$$

where $a \in \mathbb{Z}$ satisfies the condition $0 \le a < dp^N$.

Let $UD(\mathbb{Z}_p)$ be the space of uniformly differentiable functions on $\mathbb{Z}_p$. For $f \in UD(\mathbb{Z}_p)$, the p-adic q-integral on $\mathbb{Z}_p$ is defined by Kim as follows:

$$I_q(f) = \int_{\mathbb{Z}_p} f(x)d\mu_q(x) = \lim_{N\to\infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} f(x)q^x, \text{ (see [3, 4]).} \tag{4}$$

By (3) and (4), the Witt's formula for the q-Bernoulli numbers with weight α is given by

$$\int_{\mathbb{Z}_p} [x]_{q^\alpha}^n d\mu_q(x) = \widetilde{\beta}_{n,q}^{(\alpha)}, \text{ where } n \in \mathbb{Z}_+. \tag{5}$$

The q-Bernoulli polynomials with weight α are also defined by

$$\widetilde{\beta}_{n,q}^{(\alpha)}(x) = \sum_{l=0}^{n} \binom{n}{l} [x]_{q^\alpha}^{n-l} q^{\alpha l x} \widetilde{\beta}_{l,q}^{(\alpha)}. \tag{6}$$

From (4), (5) and (6), we can derive the Witt's formula for $\widetilde{\beta}_{n,q}^{(\alpha)}(x)$ as follows:

$$\int_{\mathbb{Z}_p} [x+y]_{q^\alpha}^n d\mu_q(y) = \widetilde{\beta}_{n,q}^{(\alpha)}(x), \text{ where } n \in \mathbb{Z}_+. \tag{7}$$

For $n \in \mathbb{Z}_+$ and $d \in \mathbb{N}$, the distribution relation for the q-Bernoulli polynomials with weight α are known that

$$\widetilde{\beta}_{n,q}^{(\alpha)}(x) = \frac{[d]_{q^\alpha}^n}{[d]_q} \sum_{a=0}^{d-1} q^a \widetilde{\beta}_{n,q^d}^{(\alpha)}\left(\frac{x+a}{d}\right), \text{ (see [3]).} \tag{8}$$

Recently, several authors have studied the p-adic q-Euler and Bernoulli measures on $\mathbb{Z}_p$ (see[8, 9, 11]). The purpose of this paper is to construct p-adic

q-Bernoulli distribution with weight $\alpha(= p$-adic q-Bernoulli unbounded measure with weight $\alpha)$ on $\mathbb{Z}_p$ and to study their integral representations. Finally, we construct the generalized q-Bernoulli numbers with weight α and investigate their properties related to p-adic q-L-functions.

2. p-adic q-Bernoulli distribution with weight α

Let X be any compact-open subset of $\mathbb{Q}_p$, such as $\mathbb{Z}_p$ or $\mathbb{Z}_p^*$. A p-adic distribution μ on X is defined to be an additive map from the collection of compact open set in X to $\mathbb{Q}_p$:

$$\mu\left(\bigcup_{k=1}^{n} U_k\right) = \sum_{k=1}^{n} \mu(U_k)(\text{additivity}),$$

where $\{U_1, U_2, \cdots, U_n\}$ is any collection of disjoint compact opensets in X.

The set $\mathbb{Z}_p$ has a topological basis of compact open sets of the form $a + p^n\mathbb{Z}_p$.

Consequently, if U is any compact open subset of $\mathbb{Z}_p$, it can be written as a finite disjoint union of sets

$$U = \bigcup_{j=1}^{k} (a_j + p^N\mathbb{Z}_p),$$

where $N \in \mathbb{Z}_+$ and $a_1, a_2, \cdots, a_k \in \mathbb{Z}$ with $0 \leq a_i < p^N - 1$

Indeed, the p-adic ball $a+p^n\mathbb{Z}_p$ can be represented as the union of smaller balls

$$a + p^n\mathbb{Z}_p = \bigcup_{b=0}^{p-1} (a + bp^n + p^{n+1}\mathbb{Z}_p).$$

Lemma 1. Every map μ from the collection of compact-open sets in X to $\mathbb{Q}_p$ for which

$$\mu(a + p^N\mathbb{Z}_p) = \bigcup_{b=0}^{p-1} (a + bp^N + dp^{N+1}\mathbb{Z}_p)$$

holds whenever $a + p^N\mathbb{Z}_p \subset X$, extends to a p-adic distribution on X.

Now we define a map $\mu_{k,q}^{(\alpha)}$ on the balls in $\mathbb{Z}_p$ as follows:

$$\mu_{k,q}^{(\alpha)}(a+p^n\mathbb{Z}_p)=\frac{[p^n]_{q^\alpha}^k}{[p^n]_q}q^a f_{k,q^{p^n}}^{(\alpha)}\left(\frac{\{a\}_n}{p^n}\right), \tag{9}$$

where $\{a\}_n$ is the unique number in the set $\{0,1,2,\cdots,p^n-1\}$ such that $\{a\}_n \equiv a \pmod{p^n}$.

If $a \in \{0,1,2,\cdots,p^n-1\}$, then

$$\begin{aligned}\sum_{b=0}^{p-1}\mu_{k,q}^{(\alpha)}(a+bp^n+p^{n+1}\mathbb{Z}_p)&=\sum_{b=0}^{p-1}\frac{[p^{n+1}]_{q^\alpha}^k}{[p^{n+1}]_q}q^{a+bp^n}f_{k,q^{p^{n+1}}}^{(\alpha)}\left(\frac{a+bp^n}{p^{n+1}}\right)\\&=q^a\frac{[p^n]_{q^\alpha}^k}{[p^n]_q}\frac{[p]_{(q^{p^n})^\alpha}^k}{[p]_{q^{p^n}}}\sum_{b=0}^{p-1}q^{bp^n}f_{k,(q^{p^n})^p}^{(\alpha)}\left(\frac{\frac{a}{p^n}+b}{p}\right).\end{aligned} \tag{10}$$

From (10), we note that $\mu_{k,q}^{(\alpha)}$ is p-adic distribution on $\mathbb{Z}_p$ if and only if

$$\frac{[p]_{(q^{p^n})^\alpha}^k}{[p]_{q^{p^n}}}\sum_{b=0}^{p-1}q^{bp^n}f_{k,(q^{p^n})^p}^{(\alpha)}\left(\frac{\frac{a}{p^n}+b}{p}\right)=f_{k,q^{p^n}}^{(\alpha)}\left(\frac{a}{p^n}\right).$$

Theorem 2. Let $\alpha \in \mathbb{N}$ and $k \in \mathbb{Z}_+$. Then we see that $\mu_{k,q}^{(\alpha)}(a+p^n\mathbb{Z}_p)$ is p-adic distribution on $\mathbb{Z}_p$ if and only if

$$\frac{[p]_{(q^{p^n})^\alpha}^k}{[p]_{q^{p^n}}}\sum_{b=0}^{p-1}q^{bp^n}f_{k,(q^{p^n})^p}^{(\alpha)}\left(\frac{\frac{a}{p^n}+b}{p}\right)=f_{k,q^{p^n}}^{(\alpha)}\left(\frac{a}{p^n}\right).$$

We set

$$f_{k,q^{p^n}}^{(\alpha)}(x)=\widetilde{\beta}_{k,q^{p^n}}^{(\alpha)}(x). \tag{11}$$

From (9) and (11), we get

$$\mu_{k,q}^{(\alpha)}(a+p^n\mathbb{Z}_p)=\frac{[p^n]_{q^\alpha}^k}{[p^n]_q}q^a\widetilde{\beta}_{k,q^{p^n}}^{(\alpha)}\left(\frac{a}{p^n}\right). \tag{12}$$

By (8), (12) and Theorem 2, we obtain the following theorem.

Theorem 3. Let $\mu_{k,q}^{(\alpha)}$ be given by

$$\mu_{k,q}^{(\alpha)}(a+dp^N\mathbb{Z}_p)=\frac{[dp^N]_{q^\alpha}^k}{[dp^N]_q}q^a\widetilde{\beta}_{k,q^{dp^N}}^{(\alpha)}\left(\frac{a}{dp^N}\right). \tag{13}$$

Then $\mu_{k,q}^{(\alpha)}$ extends to a $\mathbb{Q}(q)$–valued distribution on the compact open sets $U\subset X$.

From (13), we note that

$$\begin{aligned}\int_X d\mu_{k,q}^{(\alpha)}(x)&=\lim_{N\to\infty}\sum_{x=0}^{dp^N-1}\mu_{k,q}^{(\alpha)}(x+dp^N\mathbb{Z}_p)\\&=\lim_{N\to\infty}\frac{[dp^N]_{q^\alpha}^k}{[dp^N]_q}\sum_{a=0}^{dp^N-1}q^a\widetilde{\beta}_{k,q^{dp^N}}^{(\alpha)}\left(\frac{a}{dp^N}\right).\end{aligned} \tag{14}$$

By (8) and (14), we get

$$\int_X d\mu_{k,q}^{(\alpha)}(x)=\widetilde{\beta}_{k,q}^{(\alpha)}.$$

Therefore, we obtain the following theorem.

Theorem 4. For $\alpha\in\mathbb{N}$ and $k\in\mathbb{Z}_+$, we have

$$\int_X d\mu_{k,q}^{(\alpha)}(x)=\widetilde{\beta}_{k,q}^{(\alpha)}.$$

Let χ be Dirichlet character with conductor $d\in\mathbb{N}$. Then we define the generalized q-Bernoulli numbers attached to χ as follows:

$$\begin{aligned}\widetilde{\beta}_{n,\chi,q}^{(\alpha)}&=\int_X\chi(x)[x]_{q^\alpha}^n d\mu_q(x)\\&=\frac{[d]_{q^\alpha}^n}{[d]_q}\sum_{a=0}^{d-1}q^a\chi(a)\widetilde{\beta}_{n,q^d}^{(\alpha)}\left(\frac{a}{d}\right).\end{aligned} \tag{15}$$

From (13) and (15), we can derive the following equation.

$$\begin{aligned}
&\int_X \chi(x)d\mu_{k,q}^{(\alpha)}(x) = \lim_{N\to\infty}\sum_{x=0}^{dp^N-1}\chi(x)\mu_{k,q}^{(\alpha)}(x+dp^N\mathbb{Z}_p)\\
&= \lim_{N\to\infty}\frac{[dp^N]_{q^\alpha}^k}{[dp^N]_q}\sum_{x=0}^{dp^N-1}\chi(x)q^x\widetilde{\beta}_{k,q^{dp^N}}^{(\alpha)}\left(\frac{x}{dp^N}\right)\\
&= \frac{[d]_{q^\alpha}^k}{[d]_q}\sum_{a=0}^{d-1}q^a\chi(a)\left\{\lim_{N\to\infty}\frac{[p^N]_{q^{\alpha d}}^k}{[p^N]_{q^d}}\sum_{x=0}^{p^N-1}q^{dx}\widetilde{\beta}_{k,q^{dp^N}}\left(\frac{\frac{a}{d}+x}{p^N}\right)\right\}\\
&= \frac{[d]_{q^\alpha}^k}{[d]_q}\sum_{a=0}^{d-1}q^a\chi(a)\widetilde{\beta}_{k,q^d}^{(\alpha)}\left(\frac{a}{d}\right) = \widetilde{\beta}_{k,\chi,q}^{(\alpha)},
\end{aligned}$$

and

$$\begin{aligned}
&\int_{pX}\chi(x)d\mu_{k,q}^{(\alpha)}(x) = \lim_{N\to\infty}\frac{[dp^{N+1}]_{q^\alpha}^k}{[dp^{N+1}]_q}\sum_{x=0}^{dp^N-1}\chi(px)q^{px}\widetilde{\beta}_{k,q^{dp^{N+1}}}^{(\alpha)}\left(\frac{px}{dp^{N+1}}\right)\\
&= \frac{[p]_{q^\alpha}^k}{[p]_q}\frac{[d]_{q^{p\alpha}}^k}{[d]_{q^p}}\sum_{a=0}^{d-1}\chi(pa)q^{pa}\lim_{N\to\infty}\frac{[p^N]_{q^{dp\alpha}}^k}{[p^N]_{q^{dp}}}\sum_{x=0}^{p^N-1}q^{pdx}\widetilde{\beta}_{k,q^{pdp^N}}^{(\alpha)}\left(\frac{p(xd+a)}{pdp^N}\right)\\
&= \frac{[p]_{q^\alpha}^k}{[p]_q}\frac{[d]_{q^{\alpha p}}^k}{[d]_{q^p}}\sum_{a=0}^{d-1}\chi(p)\chi(a)q^{pa}\widetilde{\beta}_{k,q^{pd}}^{(\alpha)}\left(\frac{a}{d}\right) = \chi(p)\frac{[p]_{q^\alpha}^k}{[p]_q}\widetilde{\beta}_{k,\chi,q^p}^{(\alpha)}.
\end{aligned}$$

For $\beta(\neq 1)\in X^*$, we have

$$\int_{pX}\chi(x)d\mu_{k,q^{1/\beta}}^{(\alpha)}(\beta x) = \chi(\frac{p}{\beta})\frac{[p]_{q^{\alpha/\beta}}^k}{[p]_{q^{1/\beta}}}\widetilde{\beta}_{k,\chi,q^{p/\beta}}^{(\alpha)},$$

and

$$\int_X\chi(x)d\mu_{k,q^{1/\beta}}^{(\alpha)}(\beta x) = \chi(\frac{1}{\beta})\widetilde{\beta}_{k,\chi,q^{1/\beta}}^{(\alpha)}.$$

Therefore, we obtain the following theorem.

Theorem 5. For $\beta(\neq 1) \in X^*$, we have

$$\begin{aligned}
&\int_X \chi(x) d\mu^{(\alpha)}_{k,q}(x) = \widetilde{\beta}^{(\alpha)}_{k,\chi,q}, \\
&\int_{pX} \chi(x) d\mu^{(\alpha)}_{k,q}(x) = \chi(p) \frac{[p]^k_{q^\alpha}}{[p]_q} \widetilde{\beta}^{(\alpha)}_{k,\chi,q^p}, \\
&\int_{pX} \chi(x) d\mu^{(\alpha)}_{k,q^{1/\beta}}(\beta x) = \chi(\frac{p}{\beta}) \frac{[p]^k_{q^{\alpha/\beta}}}{[p]_{q^{1/\beta}}} \widetilde{\beta}^{(\alpha)}_{k,\chi,q^{p/\beta}}, \\
&\int_X \chi(x) d\mu^{(\alpha)}_{k,q^{1/\beta}}(\beta x) = \chi(\frac{1}{\beta}) \widetilde{\beta}^{(\alpha)}_{k,\chi,q^{1/\beta}}.
\end{aligned}$$

Define

$$\mu^{(\alpha)}_{k,\beta,q}(U) = \mu^{(\alpha)}_{k,q}(U) - \beta^{-1} \frac{[\beta^{-1}]^k_{q^\alpha}}{[\beta^{-1}]_q} \mu^{(\alpha)}_{k,q^{1/\beta}}(\beta U). \tag{16}$$

By simple calculation, we get

$$\begin{aligned}
&\int_{X^*} \chi(x) d\mu^{(\alpha)}_{k,\beta,q}(x) \\
&= \int_X \chi(x) d\mu^{(\alpha)}_{k,q}(x) - \beta^{-1} \frac{[\beta^{-1}]^k_{q^\alpha}}{[\beta^{-1}]_q} \int_{pX} \chi(x) \mu^{(\alpha)}_{k,q^{1/\beta}}(x) \\
&= \widetilde{\beta}^{(\alpha)}_{k,\chi,q} - \chi(p) \frac{[p]^k_{q^\alpha}}{[p]_q} \widetilde{\beta}^{(\alpha)}_{k,\chi,q^p},
\end{aligned} \tag{17}$$

and

$$\frac{[\beta^{-1}]^k_{q^\alpha}}{[\beta^{-1}]^k_q} \int_{X^*} \chi(x) d\mu^{(\alpha)}_{k,q^{1/\beta}}(\beta x) = \frac{[\frac{1}{\beta}]^k_{q^\alpha}}{[\frac{1}{\beta}]_q} \chi(1/\beta) \widetilde{\beta}^{(\alpha)}_{k,\chi,q^{1/\beta}} - \chi(p/\beta) \frac{[\frac{p}{\beta}]^k_{q^\alpha}}{[\frac{p}{\beta}]_q} \widetilde{\beta}^{(\alpha)}_{k,\chi,q^{p/\beta}}. \tag{18}$$

By (16), (17) and (18), we get

$$
\begin{aligned}
&\int_{X^*} \chi(x) d\mu^{(\alpha)}_{k,\beta,q}(\beta x) \\
&= \int_X \chi(x) d\mu^{(\alpha)}_{k,q}(x) - \beta^{-1} \frac{[\beta^{-1}]^k_{q^\alpha}}{[\beta^{-1}]_q} \int_{pX} \chi(x) \mu^{(\alpha)}_{k,q^{1/\beta}}(\beta x) \\
&= \widetilde{\beta}^{(\alpha)}_{k,\chi,q} - \chi(p) \frac{[p]^k_{q^\alpha}}{[p]_q} \widetilde{\beta}^{(\alpha)}_{k,\chi,q^p} - \frac{1}{\beta} \frac{[\frac{1}{\beta}]^k_{q^\alpha}}{[\frac{1}{\beta}]_q} \chi(1/\beta) \widetilde{\beta}^{(\alpha)}_{k,\chi,q^{1/\beta}} \\
&\qquad + \chi(p/\beta) \frac{[\frac{p}{\beta}]^k_{q^\alpha}}{[\frac{p}{\beta}]_q} \widetilde{\beta}^{(\alpha)}_{k,\chi,q^{p/\beta}}.
\end{aligned}
\tag{19}
$$

Now we define the operator $\chi^y = \chi^{y,k,\alpha:q}$ on $f(q)$ by

$$
\chi^y f(q) = \chi^{y,k,\alpha:q} f(q) = \frac{[y]^k_{q^\alpha}}{[y]_q} \chi(y) f(q^y). \tag{20}
$$

Thus, by (20), we get

$$
\begin{aligned}
&\chi^{x,k,\alpha:q} \circ \chi^{y,k,\alpha:q} f(q) = \chi^{x,k,\alpha:q} \frac{[y]^k_{q^\alpha}}{[y]_q} \chi(y) f(q^y) \\
&= \frac{[y]^k_{q^\alpha}}{[y]_q} \chi(y) \chi(x) \frac{[y]^k_{q^{\alpha y}}}{[y]_{q^y}} \chi(y) f(q^{xy}) \\
&= \frac{[xy]^k_{q^\alpha}}{[xy]_q} \chi(xy) f(q^{xy}) \\
&= \chi^{xy,k,\alpha:q} f(q) = \chi^{xy} f(q).
\end{aligned}
\tag{21}
$$

Let us define $\chi^x \chi^y = \chi^{x,k,\alpha:q} \circ \chi^{y,k,\alpha:q}$. Then we have

$$
\chi^x \chi^y = \chi^{xy}.
$$

From the definition of χ^x, we can easily derive the following equation.

$$
(1 - \chi^p) \left(1 - \frac{1}{\beta} x^{1/\beta} \right) = 1 - \frac{1}{\beta} x^{1/\beta} - \chi^p + \frac{1}{\beta} x^{p/\beta}.
$$

Let $f(q) = \widetilde{\beta}_{k,\chi,q}^{(\alpha)}$. Then we get

$$\begin{aligned}
&(1-\chi^p)\left(1-\frac{1}{\beta}x^{1/\beta}\right)\widetilde{\beta}_{k,\chi,q}^{(\alpha)} \\
&= \widetilde{\beta}_{k,\chi,q}^{(\alpha)} - \frac{1}{\beta}\frac{[\frac{1}{\beta}]_{q^\alpha}^k}{[\frac{1}{\beta}]_q}\chi(1/\beta)\widetilde{\beta}_{k,\chi,q}^{(\alpha)} - \frac{[p]_{q^\alpha}^k}{[p]_q}\chi(p)\widetilde{\beta}_{k,\chi,q^p}^{(\alpha)} \\
&\quad + \frac{1}{\beta}\frac{[\frac{p}{\beta}]_{q^\alpha}^k}{[\frac{p}{\beta}]_q}\chi(p/\beta)\widetilde{\beta}_{k,\chi,q^{p/\beta}}^{(\alpha)}.
\end{aligned} \tag{22}$$

By (19) and (22), we obtain the following equation:

$$\int_{X^*}\chi(x)d\mu_{k,\beta,q}^{(\alpha)}(\beta x) = (1-\chi^p)\left(1-\frac{1}{\beta}x^{1/\beta}\right)\widetilde{\beta}_{k,\chi,q}^{(\alpha)},$$

where $\beta(\neq 1) \in X^*$.

References

[1] L. Carlitz, Expansions of q-Bernoulli numbers. Duke Math. J. 25 (1958), 355-364

[2] L. Carlitz, q-Bernoulli numbers and polynomials. Duke Math. J. 15 (1948), 987-1000.

[3] T. Kim, On the weighted q-Bernoulli numbers and polynomials, Advanced Studies in Contemporary Mathematics 21 (2011), 207-215

[4] T. Kim, On a q-analogue of the p-adic log gamma functions and related integrals, J. Number Theory 76 (1999), 320-329

[5] Y. H. Kim, On the p-adic interpolation functions of the generalized twisted (h,q)-Euler numbers, Int. J. Math. Anal. (Ruse) 3 (2009), no. 17-20

[6] A. Bayad, Modular properties of elliptic Bernoulli and Euler functions, Adv. Stud. Contemp. Math. 20 (2010), 389–401

[7] A. Bayad, Arithmetical properties of elliptic Bernoulli and Euler numbers, Int. J. Algebra 4 (2010), 353-372.

[8] H. Ozden, Y. Simsek, S.H. Rim, I. N. Cangul , A note on p-adic q-Euler measure, Adv. Stud. Contemp. Math. 14 (2007), 233-239

[9] H. Ozden, Y. Simsek, I. N. Cangul, Euler polynomials associated with p-adic q-Euler measure, Gen. Math. 15 (2007), 24-37

[10] Y. Simsek, Special functions related to Dedekind-type DC-sums and their applications, Russ. J. Math. Phys. 17 (2010), 495-508

[11] S. H. Rim, A note on p-adic Euler measures, Russ. J. Math. Phys. 13 (2006), 358–361

[12] S.-H. Rim, J.-H. Jin, E.-J. Moon, S.-J. Lee, Some identities on the q-Genocchi polynomials of higher-order and q-Stirling numbers by the fermionic p-adic integral on $\mathbb{Z}p$, Int. J. Math. Math. Sci. 2010, Art. ID 860280, 14 pp.